

# GEMS OF TCS

## GÖDEL'S INCOMPLETENESS

---

Sasha Golovnev

October 11, 2023

# GÖDEL'S INCOMPLETENESS THEOREM



# AXIOMATIZATION OF MATH

- Find a set of simple and obvious axioms

# AXIOMATIZATION OF MATH

- Find a set of simple and obvious axioms
- Any proof could be (in principle) traced back to this set of axioms

## EUCLID'S AXIOMS

- For any pair of distinct points, there is exactly one line connecting them

## EUCLID'S AXIOMS

- For any pair of distinct points, there is exactly one line connecting them
- Any line segment can be extended to an infinite line

## EUCLID'S AXIOMS

- For any pair of distinct points, there is exactly one line connecting them
- Any line segment can be extended to an infinite line
- For any pair of distinct points, there is exactly one circle centered at the first and touching the second

## EUCLID'S AXIOMS

- For any pair of distinct points, there is exactly one line connecting them
- Any line segment can be extended to an infinite line
- For any pair of distinct points, there is exactly one circle centered at the first and touching the second
- All right angles are equal to one another



## EUCLID'S AXIOMS

- For any pair of distinct points, there is exactly one line connecting them
- Any line segment can be extended to an infinite line
- For any pair of distinct points, there is exactly one circle centered at the first and touching the second
- All right angles are equal to one another
- [The Parallel Postulate] Given a line  $L$  and a point  $x$ , there is exactly one line parallel to  $L$  that passes through  $x$

# EUCLID'S AXIOMS

- For any pair of distinct points, there is exactly one line connecting them
- Any line segment can be extended to an infinite line
- For any pair of distinct points, there is exactly one circle centered at the first and touching the second
- All right angles are equal to one another
- [The Parallel Postulate] Given a line  $L$  and a point  $x$ , there is exactly one line parallel to  $L$  that passes through  $x$

# PEANO ARITHMETIC

- 0 is a natural number

# PEANO ARITHMETIC

- 0 is a natural number
- $\forall x, x = x$
- If  $x = y$ , then  $y = x$
- If  $x = y$  and  $y = z$ , then  $x = z$
- ...

# PEANO ARITHMETIC

- 0 is a natural number
- $\forall x, x = x$
- If  $x = y$ , then  $y = x$
- If  $x = y$  and  $y = z$ , then  $x = z$
- ...
- $\forall x, y, x = y$  iff  $\text{Next}(x) = \text{Next}(y)$
- If  $x$  is a natural number, then  $\text{Next}(x)$  is a natural number
- ...

# PEANO ARITHMETIC

- 0 is a natural number
- $\forall x, x = x$
- If  $x = y$ , then  $y = x$
- If  $x = y$  and  $y = z$ , then  $x = z$
- ...
- $\forall x, y, x = y$  iff  $\text{Next}(x) = \text{Next}(y)$
- If  $x$  is a natural number, then  $\text{Next}(x)$  is a natural number
- ...
- $\forall x, y, x + \text{Next}(y) = \text{Next}(x + y)$

# PEANO ARITHMETIC

- 0 is a natural number
- $\forall x, x = x$
- If  $x = y$ , then  $y = x$
- If  $x = y$  and  $y = z$ , then  $x = z$
- ...
- $\forall x, y, x = y$  iff  $\text{Next}(x) = \text{Next}(y)$
- If  $x$  is a natural number, then  $\text{Next}(x)$  is a natural number
- ...
- $\forall x, y, x + \text{Next}(y) = \text{Next}(x + y)$
- $\forall x, y, x \cdot \text{Next}(y) = x \cdot y + x$

# PEANO ARITHMETIC

- 0 is a natural number
- $\forall x, x = x$
- If  $x = y$ , then  $y = x$
- If  $x = y$  and  $y = z$ , then  $x = z$
- ...
- $\forall x, y, x = y$  iff  $\text{Next}(x) = \text{Next}(y)$
- If  $x$  is a natural number, then  $\text{Next}(x)$  is a natural number
- ...
- $\forall x, y, x + \text{Next}(y) = \text{Next}(x + y)$
- $\forall x, y, x \cdot \text{Next}(y) = x \cdot y + x$
- Induction



# NAIVE SET THEORY

- Set
- Membership in a Set
- Empty Set
- Equality

# RUSSELL'S PARADOX

## RUSSELL'S PARADOX

The barber is the "one who shaves all those, and those only, who do not shave themselves".  
The question is, does the barber shave himself?

# PRINCIPIA MATHEMATICA

ZFC

# GÖDEL'S INCOMPLETENESS THEOREM

Any attempt to axiomatize all of mathematics is guaranteed to fail

# HALTING PROBLEM

- Function HALT is defined as follows.

# HALTING PROBLEM

- Function HALT is defined as follows.
  - The first input is algorithm A



# HALTING PROBLEM

- Function HALT is defined as follows.
  - The first input is algorithm  $A$
  - The second input is string  $x$

# HALTING PROBLEM

- Function HALT is defined as follows.
  - The first input is algorithm  $A$
  - The second input is string  $x$
  - $\text{HALT}(A, x) = 1$  if  $A$  halts on input  $x$

# HALTING PROBLEM

- Function HALT is defined as follows.
  - The first input is algorithm  $A$
  - The second input is string  $x$
  - $\text{HALT}(A, x) = 1$  if  $A$  halts on input  $x$
  - $\text{HALT}(A, x) = 0$  if  $A$  enters infinite loop on input  $x$

# HALTING PROBLEM

- Function HALT is defined as follows.
  - The first input is algorithm  $A$
  - The second input is string  $x$
  - $\text{HALT}(A, x) = 1$  if  $A$  halts on input  $x$
  - $\text{HALT}(A, x) = 0$  if  $A$  enters infinite loop on input  $x$
- HALT is undecidable (Lecture 13)